



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Yellow fever.—In the city of Guayaquil 21 new cases of yellow fever with 6 deaths.

HAWAII.

Cholera at Honolulu.

Passed Assistant Surgeon Ramus reported the occurrence of 13 cases of cholera with 10 deaths at Honolulu between February 25 and March 2. The cases have occurred in groups of 2 or 3 in widely separated localities.

A patrol service is maintained for the purpose of preventing the natives from eating shellfish taken from the possibly contaminated shallow water.

The provisions of the quarantine regulations which apply to vessels clearing from cholera-infected ports for ports of the United States are being enforced at Honolulu and inspection is being carried out at other Hawaiian ports.

Record of Plague Infection.

Last case of human plague at Honolulu occurred July 12, 1910.

The last plague-infected rat was found at Aiea, 9 miles from Honolulu, April 12, 1910.

At Hilo the last case of human plague occurred March 23, 1910. A fatal case occurred at Honokaa, 60 miles from Hilo, December 17, 1910, and 2 fatal cases were reported January 31, 1911.

The last plague-infected rat was found at Honokaa, December 20, 1910.

Passed Asst. Surg. Ramus reports in regard to plague-prevention work, February 6:

HONOLULU.

Week ended February 4, 1911.

Total rats and mongoose taken.....	753
Rats trapped.....	747
Mongoose trapped.....	6
Rats examined bacteriologically.....	781
Classification of rats trapped:	
<i>Mus alexandrinus</i>	80
<i>Mus musculus</i>	194
<i>Mus norvegicus</i>	53
<i>Mus rattus</i>	420
Average number of traps set daily.....	1,720

INDIA.

CALCUTTA—Cholera and Plague.

Acting Asst. Surg. Allan reports, January 26:

During the week ended January 7 there were at Calcutta 12 deaths from cholera and 9 from plague; in all Bengal, 1,289 cases of plague, with 1,173 deaths; in all India, 15,415 cases, with 12,671 deaths.